



ORIGINAL ARTICLE

Open
Access

The Effect of Consistent and Specialized Freestyle Wrestling Training on Upper Limb Muscle Synergy during the Double-Leg Attack Technique: A Comparison between Elite and Sub-Elite Wrestlers

Hojat Beinabaji ¹, Mansour Eslami ^{*2}, Sayed Esmail Hosseinejad ³ & Iman Esmaili Paen Afrakoti ⁴

1. Ph.D. in Sports Biomechanics, Department of Motor Behavior and Sports Biomechanics, Faculty of Physical Education and Sport Sciences, University of Mazandaran, Mazandaran, Iran
2. Associate Professor of Sports Biomechanics, Department of Motor Behavior and Sports Biomechanics, Faculty of Sports Sciences, University of Mazandaran, Mazandaran, Iran
3. Assistant Professor of Sports Biomechanics, Department of Motor Behavior and Sports Biomechanics, Faculty of Sports Sciences, University of Mazandaran, Mazandaran, Iran
4. Associate Professor of Electrical and Electronic Engineering, Department of Electrical Engineering, Faculty of Engineering and Technology, University of Mazandaran, Mazandaran, Iran

Correspondence: Mansour Eslami: Email: m.eslami@umz.ac.ir

ABSTRACT

Background: Freestyle wrestling is a complex sport that demands precise and coordinated motor control to execute techniques successfully. The Double-Leg Attack, a fundamental offensive maneuver, requires synchronized activation of multiple upper-limb and trunk muscles. Muscle synergy, or modular control of movement, reflects how the central nervous system organizes muscles into functional units to produce efficient motion. Understanding how specialized and consistent wrestling training shapes these synergies is essential for optimizing neuromuscular coordination and improving performance levels among wrestlers of different expertise.

Methods: Thirty-eight junior Iranian male freestyle wrestlers were classified into two groups based on their competitive level: elite (n = 19), successful at national or international events, and sub-elite (n = 19), active in training without major achievements. EMG signals from five unilateral upper-limb muscles were recorded during the Double-Leg Attack maneuver. Non-negative matrix factorization (NMF) was applied to extract muscle synergy composition and temporal activation patterns, enabling comparison of modular control between groups.

Results: Three distinct muscle synergies (Syn1, Syn2, Syn3) were identified in both elite and sub-elite wrestlers. Intragroup coherence in spatial structures and activation patterns indicated a consistent organizational strategy among participants. However, elite wrestlers showed significantly greater temporal activation values than sub-elite wrestlers (P = 0.000), demonstrating enhanced timing precision and coordination within the synergy framework.

Conclusions: Consistent and specialized Freestyle wrestling training appears to refine temporal control and synergy organization, contributing to superior performance. These findings underline the value of muscle synergy analysis in developing targeted training programs aimed at improving coordination and technical efficiency in wrestling.

KEY WORDS : Freestyle Wrestling, Double-Leg Attack, Motor Module Composition, Spatial Structures, Temporal Activation Patterns

<https://jast.uma.ac.ir/>

How to cite: Beinabaji, H., Eslami, M., Hosseinijad, S. Esmail, Esmaili Paeen Afrakoti, I. The Effect of Consistent and Specialized Freestyle Wrestling Training on Muscle Synergy during the Double-Leg Attack Technique: A Comparison between Elite and Sub-Elite Wrestlers. *Journal of Advanced Sport Technology*, 2025; 9(4): -. doi: 10.22098/jast.2025.4078

Introduction

Freestyle wrestling is a highly dynamic and physically demanding sport that requires precise movement coordination and rapid execution of techniques. The effectiveness of specific wrestling maneuvers, such as the Double-Leg Attack Technique, is crucial in gaining an advantageous position and securing victory. The successful application of these techniques relies not only on physical attributes like strength and agility but also on the coordinated activation of underlying muscle groups. Understanding the underlying mechanisms and adaptations associated with this technique is essential for comprehending the effects of consistent and specialized wrestling training on performance.

Motor modules, also known as muscle synergies, provide valuable insights into the neural control of movement. These structures consist of groups of muscles that work together in a specific manner, with a fixed spatial structure and a time-varying activation pattern[1]. Motor modules act as fundamental units that translate desired movements into biomechanical outputs[2–6]. They can be scaled, shifted, and combined to facilitate the coordination of intentional, goal-directed actions[7,8]. Investigating the characteristics and differences in muscle synergies among athletes of different skill levels is crucial for understanding superior performance in sports.

Long-term training can lead to structural and organizational changes in motor modules, particularly in individuals with motor impairments[9]. Studies on walking and balance abilities have shown that individuals with impairments exhibit a reduction in the number of motor modules and an increase in the number of muscles working within each module[10–13]. Furthermore, there is broader recruitment of these modules over time and a higher degree of variability in the recruitment pattern[14]. In the context of sports, studies in various disciplines have explored the relationship between skill level and muscle synergies, revealing variations in the characteristics of motor modules, including the number, spatial structure, temporal patterns, and recruitment of modules.

Studies in sports such as cycling[15], badminton[16], and beam walking[9] have demonstrated that experienced athletes exhibit a greater number of muscle synergies compared to their less experienced counterparts. However, some studies did not find significant differences in the number of muscle synergies among athletes of different skill levels[17–27], but observed variations in other characteristics or variables associated with muscle synergies. For example, novice archers showed differences in the timing and muscle contributions of specific motor modules compared to elite and mid-level archers[17]. Consistent and specialized training can potentially induce changes in the spatial and temporal structure of motor modules, with expert athletes recruiting a greater number of modules, exhibiting reduced muscle coactivity, and demonstrating more consistent motor modules during challenging sport tasks.

The objective of our study is to investigate the differences in upper limb muscle synergies between elite and sub-elite wrestlers, providing insights into the motor control mechanisms associated with superior performance. We hypothesize that consistent and specialized wrestling training alters

muscle coordination during the Double-Leg Attack maneuver. Specifically, we examine and compare the structure and organization of motor modules between elite and sub-elite wrestlers during this commonly used technique. We expect that elite wrestlers will exhibit a greater number of motor modules and demonstrate a higher level of consistency in both spatial structure and temporal patterns during the execution of the Double-Leg Attack maneuver. The findings of this study have the potential to inform training strategies, optimize performance enhancement programs, and contribute to the advancement of freestyle wrestling training methodologies.

Material and Methods

Participants

Thirty-eight junior Iranian male wrestlers without any orthopedic or neurologic abnormalities performed the Double-Leg Attack against a defender. The Elite group comprised 19 wrestlers (age: 16.7 ± 1.0 years; height: 1.73 ± 0.07 meters; mass: 65.0 ± 14.2 kilograms) who have achieved notable success in national and international competitions, while the Sub-elite group comprised 19 talented wrestlers (age: 16.8 ± 0.65 years; height: 1.75 ± 0.05 meters; mass: 67.7 ± 11.0 kilograms) who are actively engaged in training but have not yet reached the same competitive level. The sole defender was a national level wrestler (age: 17 years; height: 1.74 meters; mass: 72 kilograms) who did not take part in the study as an attacker. Prior to their participation, all participants provided written informed consent, adhering to the guidelines outlined in the Declaration of Helsinki. Furthermore, the study protocols were reviewed and approved by the ethics committee of the University of Mazandaran.

Experimental Procedures

The target task for this study was designated as the Double-Leg Attack maneuver. In the sport of freestyle wrestling, the initiation of the Double-Leg Attack involves the attacker taking a step towards the opponent using the leading foot from a staggered crouched position. Simultaneously, the attacker lowers their body by flexing the lower extremities while advancing towards the opponent. Upon reaching the desired position, the attacker forcefully propels their upper body into the abdomen of the opponent, simultaneously grasping the back of the knees and exerting a pulling force to bring the opponent down to the ground [28]. Prior to initiating each leg attack, both the attacker and the defender assumed a staggered position, with their hands in contact with their shoulders or arms, replicating the stance used in a wrestling match (see Figure 1 for reference). This approach is consistent with the methodology employed in a study conducted by Yamashita et al [29]. In order to minimize variability and improve the reliability of the collected data during this technique, we implemented a shortened movement task. Wrestlers were provided with standardized instructions to perform the technique, resulting in enhanced repeatability of their performances. Afterwards, the quality of the attacks was assessed by a wrestling expert using a rating scale ranging from 1 (poor) to 5 (excellent). Subsequently, seven trials that received a rating of at least 4 or 5 were chosen for further analysis.




Figure 1. Visual representation illustrating double-leg attack from the starting position (left) to the ending position (right).

Data Collection and Processing

Surface EMG activity during Double-Leg Attack was recorded from the five muscles of preferred upper limb of each participant using a surface EMG system (Biovision, Germany) at a sampling rate of 1000 Hz. The muscles that were recorded included the following: long head of Triceps brachii (TB), Biceps brachii (BB), Pectoralis Major (PM), Deltoid Anterior (DA), and Latissimus dorsi (LD). Bipolar Ag-AgCl electrodes (INTCO, SF06), with a diameter of 10 mm and inter-electrode distance of 20 mm, were used. To ensure minimal impedance, the skin was prepared prior to electrode application by shaving and cleaning with a mixture of alcohol and ether. Each electrode was placed longitudinally with respect to the underlying muscle fibers arrangement, adhering to the guidelines recommended by SENIAM (Surface EMG for Non-Invasive Assessment of Muscles) for all the muscles, except for the muscles latissimus dorsi (LD) and Pectoralis major (PM) which are not referenced by SENIAM. For LD and PM, the electrode placement followed the recommendations provided by Criswell [30].

The identification of the start and end of the technique in the pilot tests was facilitated by the utilization of reflective markers, three synchronized cameras (Basler, Germany) with an EMG device. The biceps muscle activity pattern, as determined by a custom-made MATLAB program, coincided with the start and end of the technique captured by the camera. Consequently, the activation pattern of the biceps muscle served as the criterion for determining the activation period of the muscles during the technique. refer to the description provided in Figure 2 for further details.




Figure 2 illustrates the determination of the onset and offset of data collection, which was achieved by leveraging the synchronous EMG and kinematic data. Specifically, we examined the resultant velocity of the wrist marker, which was plotted on the linear envelope of the EMG data. As the technique began, the resultant velocity of the wrist marker crossed the zero line and exhibited a significant increase. Conversely, at the end of the technique, the resultant velocity of the wrist marker crossed zero again, but this time displayed a significant decrease. Notably, we observed that the onset and offset of the biceps muscle coincided with the start and end of the technique, respectively. Consequently, the activation pattern of the biceps muscle served as the criterion for determining the activation period of the muscles during the technique.

Muscle Synergies Extraction

To enhance the quality of the surface electromyography (EMG) data by eliminating signal outliers without excessively impacting the raw data, a Hampel filter was employed as the initial step. The Hampel filter was configured with specific parameters, including a time window of $\text{win} = 25$ and a threshold of $\sigma = 4$ (standard deviations), to accomplish this objective [31]. Subsequently, the EMG signals underwent further processing. Firstly, a bandpass filter was applied using a 4th order Butterworth filter, restricting the frequency range between 20 and 450 Hz. Following this, the EMG data was subjected to full-wave rectification, effectively converting the signal to its absolute value, and subsequently, a 4th order zero-lag Butterworth low-pass filter was employed at 8 Hz to obtain the linear envelope. To achieve a consistent time base, the filtered data was time-interpolated with 200 points per individual cycle. To standardize the amplitude scales across subjects, the amplitudes of the EMG waveforms were normalized by dividing by the maximum value for each corresponding subject, resulting in a range of 0 to 1 for all muscle scales. Finally, the calculated data for each subject were organized into a data matrix consisting of 5 rows (representing 5 muscles) and 1400 columns (200 points per cycle multiplied by 7 cycles).

The Non-negative Matrix Factorization algorithm (NMF) was employed to extract muscle synergies from each cycle of the data matrix containing EMG recordings. The NMF approach assumes that the muscle activation pattern, denoted as M , within each period can be expressed as a linear combination of a limited number of muscle synergy composition vectors or weighing vectors, represented as W_k . Each muscle synergy is recruited by a corresponding synergy

recruitment coefficient, denoted as C_k . Thus, Equation 1 represents the representation of a specific muscle activation pattern, \mathbf{M} , in each task.

$$M(t) = \sum_{k=1}^{K_{syng}} W_k C_{k(t)} + e \quad (W_k \geq 0, C_k \geq 0) \quad \text{Equation 1}$$

where variable \mathbf{M} represents an initial matrix with dimensions m-by-n, where m corresponds to the number of muscles and n represents the number of time points. The matrix \mathbf{W} is m-by-k, where k denotes the number of synergies. \mathbf{W} represents the matrix of muscle synergy composition vectors. The matrix \mathbf{C} , with dimensions k-by-n, represents the synergy activation coefficients or temporal activation patterns. Lastly, the matrix \mathbf{e} is m-by-n and represents the residual error matrix. The NMF analysis utilized the Alternating Least Squares (ALS) algorithm, which iteratively updated initial randomly estimated matrices \mathbf{W} and \mathbf{C} . This iterative process facilitated the convergence of the matrices, leading to a locally optimal matrix factorization. The analysis applied 100 replicates and a maximum of 50 iterations, with the latter representing the default maximum value. For each subject, the analysis was performed iteratively by varying the number of synergies from 1 to 5. The aim was to identify the minimum number of synergies that accounted for more than 90% of the overall variance accounted for (VAF) and more than 75% of the VAF in each muscle. This threshold for selecting the number of synergies was consistent with the approach used in the study by Nishida et al [32]. To enable comparisons of synergy variables between individuals and groups, all muscle synergy composition vectors (\mathbf{W}) and synergy activation coefficients (\mathbf{C}) were normalized by their respective norms.

Similarity Analysis

We used cosine similarity (CS) analysis to examine the similarity in motor module composition and temporal activation patterns among wrestlers [33,34]. The cosine similarity (CS) is calculated as the inner product of two paired synergy vectors that have been normalized to unit norm. This similarity measure quantifies the cosine of the angle between the vectors and has a range from -1 to 1. Values closer to 1 indicate a higher degree of similarity between the vectors, with a value of 0 indicating orthogonality (the vectors are perpendicular to each other), and a value of -1 indicating that the vectors are diametrically opposed or have opposite directions. Since the synergy vectors used in this context are nonnegative, the computation of cosine similarity (CS) yields values ranging from 0 to 1. The CS was calculated for each pair of wrestlers by comparing their respective synergy vectors. Two vectors with $CS > 0.8$ were defined as similar [35,36].

Statistical Analysis

The Shapiro-Wilk test was utilized to evaluate the normal distribution of the data. To compare the evolution of variance accounted for (VAF) with the number of extracted synergies between the two groups, an analysis of variance (ANOVA) for repeated measures was employed [27]. An Independent-Samples T-Test was used to compare characteristics related to muscle weighing and coefficient activation synergies between the two groups. When the distribution of data was not normal, Mann-Whitney U test was used. All statistical analyses were performed with SPSS version 27. A $p \leq 0.05$ was considered indicative of statistical significance. All analyses were conducted using custom software developed in MATLAB R2023a.

Results

Number of muscle synergies: A non-significant difference ($p = 0.133$) was observed between the two groups regarding the evolution of VAF as a function of the number of muscle synergies. This finding indicates that both groups exhibit similar dimensionality in their EMG data. The results are illustrated in Figure 3 (left). Applying the previously described criteria, muscle synergies ranging from one to four were identified for all conditions and groups. However, in most conditions, the EMG activity was adequately explained by three synergies. The distribution of variation appeared to follow a normal distribution, as depicted in Figure 3 (right). Based on the observation that three synergies adequately accounted for the EMG activity in all conditions, a conclusive number of synergies was set to three. Subsequently, the NMF algorithm was again applied to all conditions using this new number of synergies. The three synergies were arranged in a specific order based on the timing of the main peak of their activation patterns. Following this order, the synergies were named Syn1, Syn2, and Syn3.




Figure 3. A: The number of motor modules selected accounted for >90% of the overall variability accounted for (VAF) and >75 in each muscle. There was no statistically significant difference in the number of muscle synergies between the groups. **B:** Histogram of the number of synergies determined by the variance accounted for (VAF) threshold criterion. The histogram includes trials for all subjects.




Figure 4. A: Synergy Activation Coefficient or Temporal Activation Patterns of three motor modules during throwing double leg attack. (mean over cycles \pm SD). Blue and red represent Elite and Sub-elite wrestlers respectively. **B:** Representative of muscle weighting synergies extracted from elite and sub-elite wrestlers. The values for muscle weighting are presented as mean \pm standard deviation (SD). Asterisks (*) indicate significant differences at $p < 0.05$. Blue and red represent Elite and Sub-elite wrestlers respectively.

Temporal Activation Patterns (Synergy Activation Coefficients): The synergy activation coefficients for elite and sub-elite wrestlers were depicted in Figure 4A. In both groups the activity of the three temporal activation patterns peaked in sequential order. Cosine similarity (CS) analysis was employed to assess the consistency of synergy activation coefficients. The heat map in Figure 5A displays similarity in temporal activation patterns across all extracted synergies from all wrestlers. Elite wrestlers demonstrated a notable degree of intragroup similarity across all three temporal activation patterns with a median CS value greater than 0.80, indicating a high level of consistency. In contrast, sub-elite wrestlers exhibited moderate to high similarity within these temporal activation patterns, which was significantly lower compared to the observed similarity in elite wrestlers. (See table 1: A and Figure 6A)




Figure 5. The heat maps in Figure A and B illustrate the interindividual similarity in the synergy activation coefficients and motor module compositions, respectively, among all wrestlers. Each cell in the heat map represents the cosine similarity for a pair of wrestlers, with the diagonal cells representing self-comparisons that are maximally similar. These heat maps provide a visual representation of the similarities between wrestlers in terms of their synergy activation coefficients and motor module compositions.




Figure 6. presents boxplots showcasing the distribution of cosine similarity values between each pair of players within three synergies for both elite and sub-elite wrestlers. Panel A compares the interindividual similarity of synergy activation coefficients, while Panel B compares the interindividual similarity of motor module composition. Significant differences between modules are indicated by their respective values.

Motor Module Composition (Muscle synergies weightings): Figure. 3B depicts the motor module composition or spatial structure of the three identified muscle synergies. In Syn1, the primary muscle is the Deltoid anterior, while in Syn3, it is the Latissimus dorsi. Notably, no

dominant muscle was observed in Syn2. Additionally, Pectoralis major, Biceps and Triceps brachii exhibited varying degrees of activity across all three synergies.

The heat map in Figure 5B displays similarity in motor module composition across all extracted synergies from all wrestlers. The motor module composition exhibited a high degree of similarity among wrestlers, with a median CS value greater than 0.80. However, for Syn3, sub-elite wrestlers displayed a higher CS value. Overall, no significant differences were found between the groups in terms of motor module composition. (See table 1: B and Figure 6B)

Table 1: Similarity indices (median \pm IQR) for each muscle synergy within and between groups.

A: Synergies Activation Coefficients (C)	Intra-group similarity		Inter-group similarity	
	Elites	Sub-elites	P	
Synergy #1	0.91 \pm 0.11	0.80 \pm 0.21	0.000*	0.81 \pm 0.14
Synergy #2	0.89 \pm 0.08	0.82 \pm 0.13	0.000*	0.84 \pm 0.09
Synergy #3	0.90 \pm 0.08	0.87 \pm 0.10	0.000*	0.84 \pm 0.09
B: Motor Module composition (W)				
Synergy #1	0.85 \pm 0.22	0.80 \pm 0.20	0.223	0.78 \pm 0.17
Synergy #2	0.86 \pm 0.27	0.83 \pm 0.14	0.723	0.80 \pm 0.13
Synergy #3	0.81 \pm 0.33	0.86 \pm 0.16	0.002*	0.80 \pm 0.17

Discussion

In this study, we provide evidence of the impact of years of specialized wrestling training on the structural modifications of muscle coordination patterns, particularly in a commonly employed technique such as Double-Leg Attack. Significant differences observed in the temporal activation patterns suggest that elite wrestlers possess a higher level of temporal precision and coordination, which may contribute to their competitive advantage.

Number of Motor Modules: In line with the findings of several other studies [17–27], our study did not reveal a significant difference in the number of synergies between elite and sub-elite groups. However, a few studies conducted in various sports disciplines, such as cycling [15], badminton [16], and beam walking [9], have demonstrated that experienced athletes exhibit a greater number of muscle synergies compared to their less experienced counterparts. Furthermore, it has been shown that motor training during rehabilitation can lead to an increase in the number of motor modules in individuals with sensorimotor deficits [37]. The variability in the number of muscle synergies observed between different groups of expertise in various studies can be

attributed to the specific characteristics of the task or sport, participant characteristics, training duration, methodological differences, and individual variability in movement patterns.

Sawers et al [9] found that experts in beam walking have more motor modules, which provides them with more flexibility in achieving motor goals. This expanded repertoire allows experts to produce additional biomechanical functions that novices cannot, particularly in maintaining balance on the beam. In contrast to beam walking that is a challenging balance behavior, it seems Double-Leg Attack maneuver characterized by its fast and explosive nature. Therefore, this maneuver is more consistent with constrained athletic activities requiring high power output such as rowing [24,27], weightlifting [18,19], and powerlifting [26]. It is important to acknowledge that in our study, the sub-elite participants were actively involved in training, although they had not yet attained the same level of competitiveness as the elites. It is also worth mentioning that they were not inexperienced individuals. Therefore, the expectation of both groups having an equal number of modules was not far from being met.

Spatial and Temporal Characteristics of Muscle Synergies: Our study revealed that all wrestlers demonstrated a high level of consistency in both the composition and temporal activation patterns of their muscle synergies, as indicated by cosine similarity (CS) values exceeding 0.8. However, notable differences between the two groups were observed solely in the temporal activation patterns. Specifically, elite wrestlers exhibited a significantly higher level of temporal consistency in their muscle activity structure during the exercise, highlighting their superior control and coordination.

In a study conducted by Turpin et al [27], both experienced and untrained rowers displayed moderate within-group similarity in the weighting of muscle synergies (ranging from 0.64 to 0.75). However, when analyzing synergy activation patterns, both groups exhibited higher levels of intra-group similarity. Specifically, the experienced group had significantly higher similarity values for the first synergy and lower values for the third synergy compared to the untrained group. Based on these findings, the authors concluded that expertise in rowing is more likely to be linked to adjustments in the mechanical output of muscle synergies rather than differences in the shape and timing of their activations. In another study conducted by Santos et al [19], the researchers examined the muscle synergy of weightlifters and untrained individuals during the execution of the power clean exercise. Intra-group variability for each group was assessed, and it was found that the muscle synergy vectors showed weak to moderate correlations (ranging from 0.34 to 0.52) in both groups. However, the intra-group variability values in the synergy activation coefficients were high (ranging from 0.77 to 0.89) and displayed strong correlations. Interestingly, when comparing the two groups, it was observed that expertise did not significantly impact the synergistic organization of muscle coordination. Overall, when comparing these findings, our study highlights the high consistency and coordination in muscle synergies among elites in the Double-Leg Attack exercise, while the other studies shed light on the role of expertise in rowing and weightlifting regarding adjustments in muscle synergies' mechanical output and the lack of significant impact on synergistic organization across different expertise levels.

The findings of this study have several potential applications for coaches and athletes in the field of wrestling and other sports. We found that consistent and specialized training can influence muscle coordination patterns to optimize training programs. Coaches can use the muscle coordination patterns of elite wrestlers as a reference point for technique refinement. By improving the temporal and spatial consistency of muscle activity, coaches can help athletes execute wrestling maneuvers more precisely and efficiently. Additionally, the differences in muscle coordination patterns between elite and sub-elite wrestlers can be valuable for talent identification purposes,

allowing coaches and talent scouts to identify athletes with the potential for elite performance and provide them with appropriate training and support.

During this project, two important limitations arose because of the nature of wrestling techniques involving close grappling between two wrestlers and limitations in laboratory equipment availability. One limitation of this study was the absence of kinematic and kinetic data. The synchronous collection of EMG and kinematic data could enhance our understanding of movement analysis. Additionally, the study was limited by the inclusion of a low number of selected muscles from unilateral upper limb. A more extensive selection of muscles from either the ipsilateral or contralateral limb would have provided a more comprehensive and informative analysis of muscle synergies.

Conclusion

This study investigated muscle coordination patterns during the Double-Leg Attack maneuver in elite and sub-elite wrestlers. The number of muscle synergies was similar between the two groups. However, notable differences were observed in the temporal consistency of muscle activity. Elite wrestlers demonstrated a high level of consistency, indicating precise and coordinated execution of the maneuver, while sub-elite wrestlers exhibited greater interindividual variability, suggesting less precise and coordinated execution and a potential need for further refinement in their muscle coordination. These findings suggest that consistent and specialized training can influence the structure of muscle coordination patterns during wrestling techniques. Overall, this study provides valuable insights that can inform coaching strategies, training programs, performance assessment, technique refinement, injury prevention, and talent identification in the field of wrestling.

Ethical Considerations:

Compliance with ethical guidelines

The research protocol was reviewed and approved by the Ethics Committee of the University of Mazandaran (Approval Code: IR.UMZ.REC.1402.033), in full compliance with institutional ethical guidelines.

Funding

This research received no external funding.

Conflict of Interest

The authors declare that there are no conflicts of interest regarding the publication of this manuscript

Acknowledgment

We thank the young wrestlers and their coaches who participated in this study. We would also like to appreciate the full cooperation of the university officials.

References

1. Ting LH, Torres-Oviedo G. Muscle Synergies Simplifying Neural Control of Posture. *FASEB J* [Internet]. 2007 Apr;21(5). Available from: <https://onlinelibrary.wiley.com/doi/10.1096/fasebj.21.5.A463-a>
2. Neptune RR, Clark DJ, Kautz SA. Modular control of human walking: A simulation study. *J Biomech* [Internet]. 2009;42(9):1282–7. Available from: <https://www.sciencedirect.com/science/article/pii/S0021929009001389>
3. Chvatal SA, Torres-Oviedo G, Safavynia SA, Ting LH. Common muscle synergies for control of center of mass and force in nonstepping and stepping postural behaviors. *J Neurophysiol*. 2011;106(2):999–1015.
4. Chvatal SA, Ting LH. Voluntary and reactive recruitment of locomotor muscle synergies during perturbed walking. *J Neurosci*. 2012;32(35):12237–50.
5. Chvatal SA, Ting LH. Common muscle synergies for balance and walking. *Front Comput Neurosci*. 2013;7:48.
6. Hayes HB, Chvatal SA, French MA, Ting LH, Trumbower RD. Neuromuscular constraints on muscle coordination during overground walking in persons with chronic incomplete spinal cord injury. *Clin Neurophysiol*. 2014;125(10):2024–35.
7. Ting LH, Chiel HJ, Trumbower RD, Allen JL, McKay JL, Hackney ME, et al. Neuromechanical Principles Underlying Movement Modularity and Their Implications for Rehabilitation. *Neuron* [Internet]. 2015 Apr;86(1):38–54. Available from: <http://dx.doi.org/10.1016/j.neuron.2015.02.042>
8. Flash T, Hochner B. Motor primitives in vertebrates and invertebrates. *Curr Opin Neurobiol*. 2005;15(6):660–6.
9. Sawers A, Allen JL, Ting LH. Long-term training modifies the modular structure and organization of walking balance control. *J Neurophysiol* [Internet]. 2015 Dec 1;114(6):3359–73. Available from: <https://www.physiology.org/doi/10.1152/jn.00758.2015>
10. Rodriguez KL, Roemich RT, Cam B, Fregly BJ, Hass CJ. Persons with Parkinson’s disease exhibit decreased neuromuscular complexity during gait. *Clin Neurophysiol*. 2013;124(7):1390–7.
11. Fox EJ, Tester NJ, Kautz SA, Howland DR, Clark DJ, Garvan C, et al. Modular control of varied locomotor tasks in children with incomplete spinal cord injuries. *J Neurophysiol*. 2013;110(6):1415–25.
12. Clark DJ, Ting LH, Zajac FE, Neptune RR, Kautz SA. Merging of Healthy Motor Modules Predicts Reduced Locomotor Performance and Muscle Coordination Complexity Post-Stroke. *J Neurophysiol* [Internet]. 2010 Feb;103(2):844–57. Available from: <https://www.physiology.org/doi/10.1152/jn.00825.2009>
13. Gizzi L, Nielsen JF, Felici F, Ivanenko YP, Farina D. Impulses of activation but not motor modules are preserved in the locomotion of subacute stroke patients. *J Neurophysiol*. 2011;106(1):202–10.
14. Routson RL, Kautz SA, Neptune RR. Modular organization across changing task demands in healthy and poststroke gait. *Physiol Rep* [Internet]. 2014 Jun;2(6):e12055. Available from: <http://doi.wiley.com/10.14814/phy2.12055>
15. Sato T, Kurematsu R, Shigetome S, Matsumoto T, Tsuruda K, Tokuyasu T. Motor control mechanism underlying pedaling skills: an analysis of bilateral coordination in the lower extremities. *Artif Life Robot* [Internet]. 2020 May 29;25(2):308–15. Available from: <https://doi.org/10.1007/s10015-019-00580-8>
16. Matsunaga N, Kaneoka K. Comparison of Modular Control during Smash Shot between Advanced and Beginner Badminton Players. *Appl Bionics Biomech* [Internet]. 2018 Jun 13;2018:1–6. Available from: <https://www.hindawi.com/journals/abb/2018/6592357/>
17. Baifa Z, Xinglong Z, Dongmei L. Muscle coordination during archery shooting: A comparison of archers with different skill levels. *Eur J Sport Sci* [Internet]. 2023 Jan 2;23(1):54–61. Available from: <https://www.tandfonline.com/doi/full/10.1080/17461391.2021.2014573>

18. Hakariya N, Kibushi B, Okada J. Differences in muscle synergies between skilled and unskilled athletes in power clean exercise at various loads. *J Sports Sci* [Internet]. 2023 Jun 3;41(11):1136–45. Available from: <https://www.tandfonline.com/doi/full/10.1080/02640414.2023.2259268>
19. Santos PDG, Vaz JR, Correia PF, Valamatos MJ, Veloso AP, Pezarat-Correia P. Intermuscular Coordination in the Power Clean Exercise: Comparison between Olympic Weightlifters and Untrained Individuals—A Preliminary Study. *Sensors* [Internet]. 2021 Mar 9;21(5):1904. Available from: <https://www.mdpi.com/1424-8220/21/5/1904>
20. Jeon Y, Kim Y, Ji Y. Specific muscle coordination patterns of taekwondo roundhouse kick in athletes and non-athletes. *Arch BUDO*. 2021;17:135–44.
21. Rahatabad FN, Maghooli K, Balan TR. Relationship between Muscle Synergies and Skills of Basketball Players. *Iran J Med Phys*. 2021;18(1):30–9.
22. Kim M, Kim Y, Kim H, Yoon B. Specific muscle synergies in national elite female ice hockey players in response to unexpected external perturbation. *J Sports Sci* [Internet]. 2018 Feb 18;36(3):319–25. Available from: <https://www.tandfonline.com/doi/full/10.1080/02640414.2017.1306090>
23. Barnamehei H, Tabatabai Ghomsheh F, Safar Cherati A, Pouladian M. Upper Limb Neuromuscular Activities and Synergies Comparison between Elite and Nonelite Athletics in Badminton Overhead Forehand Smash. *Appl Bionics Biomech* [Internet]. 2018 Dec 20;2018:1–10. Available from: <https://www.hindawi.com/journals/abb/2018/6067807/>
24. Shaharudin S, Agrawal S. Muscle synergies during incremental rowing VO₂max test of collegiate rowers and untrained subjects. *J Sports Med Phys Fitness* [Internet]. 2016 Sep;56(9):980–9. Available from: www.researchgate.net/publication/
25. Vaz JR, Olstad BH, Cabri J, Kjendlie P-L, Pezarat-Correia P, Hug F. Muscle coordination during breaststroke swimming: Comparison between elite swimmers and beginners. *J Sports Sci* [Internet]. 2016 Oct 17;34(20):1941–8. Available from: <http://www.tandfonline.com/doi/full/10.1080/02640414.2016.1143109>
26. Kristiansen M, Madeleine P, Hansen EA, Samani A. Inter-subject variability of muscle synergies during bench press in power lifters and untrained individuals. *Scand J Med Sci Sports* [Internet]. 2015 Feb 24;25(1):89–97. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/sms.12167>
27. Turpin NA, Guével A, Durand S, Hug F. No evidence of expertise-related changes in muscle synergies during rowing. *J Electromyogr Kinesiol* [Internet]. 2011 Dec;21(6):1030–40. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1050641111001106>
28. Mysnyk M, Davis B, Simpson B. *Winning Wrestling Moves*. Human kinetics; 1994. 208 p.
29. Yamashita D, Arakawa H, Wada T, Yumoto K, Fujiyama K, Nagami T, et al. Whole-Body Mechanics of Double-Leg Attack in Elite and Non-elite Male Freestyle Wrestlers. *Front Sport Act Living* [Internet]. 2020 Jun 10;2:58. Available from: <https://www.frontiersin.org/article/10.3389/fspor.2020.00058/full>
30. Criswell E. Electrode Placements. In: *cram's introduction to surface electromyography*. 2nd ed. Jones & Bartlett Learning; 2010.
31. Bhowmik S, Jelfs B, Arjunan SP, Kumar DK. Outlier removal in facial surface electromyography through Hampel filtering technique. In: 2017 IEEE Life Sciences Conference (LSC) [Internet]. IEEE; 2017. p. 258–61. Available from: <http://ieeexplore.ieee.org/document/8268192/>
32. Nishida K, Hagio S, Kibushi B, Moritani T, Kouzaki M. Comparison of muscle synergies for running between different foot strike patterns. *Alway SE*, editor. *PLoS One* [Internet]. 2017 Feb 3;12(2):e0171535. Available from: <https://dx.plos.org/10.1371/journal.pone.0171535>
33. Hagio S, Kouzaki M. Action direction of muscle synergies in three-dimensional force space. *Front Bioeng Biotechnol*. 2015;3:187.
34. Aoyama T, Ae K, Kohno Y. Interindividual differences in upper limb muscle synergies during baseball throwing motion in male college baseball players. *J Biomech* [Internet]. 2022 Dec;145:111384. Available from: <https://doi.org/10.1016/j.jbiomech.2022.111384>
35. Oliveira AS, Silva PB, Lund ME, Gizzi L, Farina D, Kersting UG. Effects of perturbations to balance on neuromechanics of fast changes in direction during locomotion. *PLoS One*. 2013;8(3):e59029.

36. Oliveira AS, Gizzi L, Farina D, Kersting UG. Motor modules of human locomotion: influence of EMG averaging, concatenation, and number of step cycles. *Front Hum Neurosci.* 2014;8:335.
37. Routson RL, Clark DJ, Bowden MG, Kautz SA, Neptune RR. The influence of locomotor rehabilitation on module quality and post-stroke hemiparetic walking performance. *Gait Posture [Internet].* 2013 Jul;38(3):511–7. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3624763/pdf/nihms412728.pdf>

تأثیر تمرین تخصصی و مستمر کشتی آزاد بر هم‌افزایی عضلانی اندام فوقانی حین اجرای فن زیر دوخم: مقایسه میان کشتی‌گیران نخبه و معمولی

نام حجت بیناباجی^۱، منصور اسلامی^{۲*}، سید اسماعیل حسینی‌نژاد^۳ و ایمان اسمعیلی پاپین افراکتی^۴

- ۱- فارغ‌التحصیل بیومکانیک ورزشی، گروه رفتار حرکتی و بیومکانیک ورزشی، دانشکده علوم ورزشی، دانشگاه مازندران، مازندران، ایران
- ۲- دانشیار بیومکانیک ورزشی، گروه رفتار حرکتی و بیومکانیک ورزشی، دانشکده علوم ورزشی، دانشگاه مازندران، مازندران، ایران
- ۳- استادیار بیومکانیک ورزشی، گروه رفتار حرکتی و بیومکانیک ورزشی، دانشکده علوم ورزشی، دانشگاه مازندران، مازندران، ایران
- ۴- دانشیار مهندسی برق و الکترونیک، گروه مهندسی برق، دانشکده فنی و مهندسی، دانشگاه مازندران، مازندران، ایران

نویسنده مسئول: منصور اسلامی: رایانامه: m.eslami@umz.ac.ir

چکیده

هدف: اجرای فنون در کشتی آزاد نیازمند زمان‌بندی و هماهنگی حرکتی دقیق است. حمله زیر دوخم یکی از مانورهای حیاتی برای کسب امتیاز و پیروزی در مسابقه محسوب می‌شود. تحلیل هماهنگی فعالیت‌های عضلانی حین اجرای فنون ورزشی نقش مهمی در بهینه‌سازی عملکرد و بررسی مکانیزم‌ها و اثرات زیربنایی تمرینات تخصصی دارد. این مطالعه با هدف بررسی تأثیر تمرینات مداوم و تخصصی کشتی آزاد بر ساختار هم‌افزایی‌های عضلانی در زمان اجرای مانور حمله زیر دوخم انجام شد. روش‌شناسی: در این پژوهش ۳۸ کشتی‌گیر جوان مازندرانی بر اساس سطح مهارت و سوابق قهرمانی به دو گروه نخبه (۱۹ نفر) و معمولی (۱۹ نفر) تقسیم شدند. فعالیت الکترومایوگرافی (EMG) پنج عضله اندام فوقانی در سمت برتر بدن ورزشکاران ثبت شد. برای استخراج هم‌افزایی‌های عضلانی و تحلیل الگوهای فعال‌سازی زمانی از الگوریتم تجزیه ماتریس غیرمنفی (NMF) استفاده گردید.

نتایج: سه هم‌افزایی عضلانی برای هر دو گروه نخبه و معمولی استخراج شد. هم‌افزایی‌های شناسایی‌شده از نظر ساختارهای فضایی و الگوهای فعال‌سازی زمانی از ثبات بالایی برخوردار بودند. کشتی‌گیران نخبه نسبت به کشتی‌گیران معمولی، ثبات زمانی و دقت بالاتری در الگوهای فعال‌سازی عضلانی نشان دادند، که این تفاوت از نظر آماری معنی‌دار بود. نتیجه‌گیری: یافته‌های این مطالعه اهمیت تمرینات تخصصی و مستمر کشتی آزاد را در بهینه‌سازی هم‌افزایی عضلانی هنگام اجرای فن زیر دوخم تأیید می‌نماید. تفاوت‌های مشاهده‌شده در الگوهای زمانی بیانگر آن است که کشتی‌گیران نخبه از هماهنگی عصبی عضلانی دقیق‌تر و کارآمدتری برخوردارند، که می‌تواند به مزیت رقابتی آنان بینجامد. این یافته‌ها راهنمایی ارزشمند برای مربیان و ورزشکاران در طراحی برنامه‌های تمرینی هدفمند با تمرکز بر ارتقای هماهنگی عضلانی و بهبود عملکرد در کشتی آزاد ارائه می‌دهد.

واژه‌های کلیدی: کشتی آزاد، فن زیر دوخم، ترکیب مازول حرکتی، ساختارهای فضایی، الگوهای فعال‌سازی زمانی

